

ORIGINAL ARTICLE

Efficacy of classification-based cognitive functional therapy in patients with non-specific chronic low back pain: A randomized controlled trial

K. Vibe Fersum¹, P. O'Sullivan², J.S. Skouen^{1,3}, A. Smith², A. Kvåle¹

¹ Physiotherapy Research Group, Department of Public Health and Primary Health Care, University of Bergen, Norway

² School of Physiotherapy, Curtin University, Bentley, Western Australia, Australia

³ The Outpatient Spine Clinic, Department of Physical Medicine and Rehabilitation, Haukeland University Hospital, Bergen, Norway

Correspondence

Kjartan Vibe Fersum
E-mail: kjartan.fersum@isf.uib.no

Funding sources

The Norwegian Fund for Post-Graduate Training in Physiotherapy supported this study.

Conflict of interest

We declare no conflict of interest.

Accepted for publication

31 October 2012

doi:10.1002/j.1532-2149.2012.00252.x

Abstract

Background: Non-specific chronic low back pain disorders have been proven resistant to change, and there is still a lack of clear evidence for one specific treatment intervention being superior to another.

Methods: This randomized controlled trial aimed to investigate the efficacy of a behavioural approach to management, classification-based cognitive functional therapy, compared with traditional manual therapy and exercise. Linear mixed models were used to estimate the group differences in treatment effects. Primary outcomes at 12-month follow-up were Oswestry Disability Index and pain intensity, measured with numeric rating scale. Inclusion criteria were as follows: age between 18 and 65 years, diagnosed with non-specific chronic low back pain for >3 months, localized pain from T12 to gluteal folds, provoked with postures, movement and activities. Oswestry Disability Index had to be >14% and pain intensity last 14 days >2/10. A total of 121 patients were randomized to either classification-based cognitive functional therapy group ($n = 62$) or manual therapy and exercise group ($n = 59$).

Results: The classification-based cognitive functional therapy group displayed significantly superior outcomes to the manual therapy and exercise group, both statistically ($p < 0.001$) and clinically. For Oswestry Disability Index, the classification-based cognitive functional therapy group improved by 13.7 points, and the manual therapy and exercise group by 5.5 points. For pain intensity, the classification-based cognitive functional therapy improved by 3.2 points, and the manual therapy and exercise group by 1.5 points.

Conclusions: The classification-based cognitive functional therapy produced superior outcomes for non-specific chronic low back pain compared with traditional manual therapy and exercise.

1. Introduction

The current evidence for management of non-specific chronic low back pain (NSCLBP) reveals that interventions such as manual therapy, exercise, acupuncture, spinal injections and cognitive behavioural therapy are not superior to each other and have a limited long-term impact on the disorder (Assendelft

et al., 2004; Furlan et al., 2005; Hayden et al., 2005; Ostelo et al., 2005; Staal et al., 2008).

Possible reasons for the failure of current clinical practice to effectively manage NSCLBP are proposed to lie in two main domains:

(1) The failure to adequately deal with NSCLBP within a multidimensional biopsychosocial framework (Borkan et al., 2002). It has been proposed that

What's already known about this topic?

- Effect sizes from randomized controlled trials utilizing conservative treatment for non-specific chronic low back pain are small.
- Stratified treatment seems to be warranted and has socio-economic impact.
- For non-specific chronic low back pain, there does not seem to be a clear consensus as to which classification system to use.

What does this study add?

- First trial for non-specific chronic low back pain utilizing a novel multidimensional classification system and cognitive functional approach.
- Cognitive functional approach shows clinical and statistically significant large effect sizes for primary outcomes as well as for secondary outcomes across multiple dimensions.

NSCLBP represents a vicious cycle associated with different combinations of provocative factors. These include cognitive factors (such as negative beliefs, fear-avoidance behaviours, catastrophizing, hypervigilance, anxiety, depression, stress, poor pacing and maladaptive coping) (Vlaeyen and Crombez, 1999; Linton, 2000), physical factors (pain provocative postures and movement patterns related to altered body schema, muscle guarding, pain behaviours and deconditioning) (O'Sullivan et al. 2006b) and lifestyle factors (sedentary behaviour, inactivity and sleep deficits) (Bjorck-van Dijken et al., 2008).

(2) Lack of a multidimensional classification system (MDCS) directing person-centred targeted management for this large group of NSCLBP patients. The Cochrane Back Review Group proposed that identification of subgroups is a key priority of low back pain (LBP) management in order to deal with the problem of patient heterogeneity (Bouter et al., 2003). Recent research supports this claim, with evidence that NSCLBP subjects can be broadly classified based on psychological factors (Turk, 2005; Boersma and Linton, 2006; Hill et al., 2010), movement and postural behaviours (Dankaerts et al., 2009), neurophysiological factors (Smart et al., 2011) and lifestyle behaviours [sedentary (Bjorck-van Dijken et al., 2008) vs. excessive activity (Mitchell et al., 2010)]. In spite of this knowledge, a recent systematic review concluded that few clinical trials exist utilizing MDCS or targeted interventions for NSCLBP (Fersum et al., 2010). The few trials that have taken a targeted biopsychosocial approach to the management of NSCLBP have dem-

onstrated a tendency for improved outcomes (Klamer Moffett et al., 2004; Vollenbroek-Hutten et al., 2004; Asenlof et al., 2005; Riipinen et al., 2005).

A novel MDCS for LBP has been developed incorporating the biopsychosocial model (O'Sullivan, 2005). This system is integrated within the Quebec classification system (Spitzer, 1987) and represents a multilevel patient-centred clinical reasoning approach to broadly classify and target management for patients with NSCLBP. This person-centred intervention is called classification-based 'cognitive functional therapy' (CB-CFT) as it directly challenges these behaviours in a cognitively integrated, functionally specific and graduated manner. This MDCS has good inter-tester reliability (Dankaerts et al., 2006; Fersum et al., 2009), with a number of studies supporting the validity of the different subgroups on the physical domains (O'Sullivan et al., 2006a,b; O'Sullivan and Beales, 2007a,b; Beales et al., 2009; Dankaerts et al., 2009), as well as cognitive domains (Boersma and Linton, 2006). This MDCS has not been formally tested in a randomized controlled trial (RCT) for NSCLBP disorders. Therefore, the aim of this study was to investigate the efficacy of CB-CFT compared with manual therapy and exercise (MT-EX) for the management of NSCLBP. The hypothesis was that a person-centred classification-based cognitive functional approach to management of NSCLBP disorders would be more effective than MT-EX approach.

2. Methods

2.1 Study design

The study was a RCT designed to evaluate the efficacy of the CB-CFT according to the system proposed by O'Sullivan (2005), as compared with MT-EX in patients with NSCLBP. The effects of the intervention were assessed at 3- and 12-month follow-up. The study received ethical approval by the regional ethics committee of medical research in Western Norway (REK Vest). The clinical trial's register number is NCT01129817.

2.2 Participants and procedures

A Norwegian university town, Bergen, with 250,000 inhabitants, provided the setting for the study. The patients were recruited from March 2006 to June 2008 from private physiotherapy outpatient practices, general practitioners and the outpatient spine clinic at the Haukeland University Hospital. In addition, six advertisements were placed in the local newspaper. All

patients were presented with written information about the study with its aims and procedures. Here, it was clearly stated that there were two active comparable treatment arms, and that based on current knowledge, we did know which was superior. The patients gave a written informed consent prior to proceeding to the clinical examination. The participants were eligible for the study if they were between the ages of 18 and 65 years, were diagnosed with NSLBP for >3 months that was primarily localized from T12 to gluteal folds, and they reported that their pain was provoked and relieved with postures, movement and activities. Pain intensity measured with a pain intensity numerical rating scale (PINRS) over the last 14 days >2/10 and an Oswestry Disability Index (ODI) >14% was necessary to be admitted to the study.

The inclusion criteria of localized back pain with mechanical behaviour to the pain were designed to include patients whose movement behaviours had a clear association with their pain disorder, for which the CB-CFT intervention was designed.

The exclusion criteria were continuous sick-leave duration for >4 months, as it was considered that specific work-related intervention would have been required, acute exacerbation of LBP at time of testing in order to avoid regression to the mean, specific LBP diagnosis (radicular pain, disc herniation, spondylolisthesis, stenosis, Modic changes), any low limb surgery in the last 3 months, surgery involving the lumbar spine, pregnancy, diagnosed psychiatric disorder, widespread constant non-specific pain disorder, pain without a clear mechanical behaviour, active rheumatologic disease, progressive neurological disease, serious cardiac or other internal medical condition, malignant diseases, acute traumas, infections or acute vascular catastrophes. A lack of compliance of greater than 50% was set as a withdrawal criterion based on programme compliance forms. An a priori power calculation suggested that 63 patients in each group would give 80% power to detect a group difference of 5 points in the ODI [estimating a standard deviation (SD) of 10 points in both groups], and 1 point on the PINRS at 1-year post-intervention (estimating an SD of 2 points in both group).

The randomization took place at the University of Bergen (UiB). A person independent of the study developed a randomization schedule and produced 160 sealed opaque envelopes containing each participant's allocation. Randomization was performed in permuted blocks of 16. When the patients had been examined and classified and a blinded examiner had collected baseline data, the patient drew the envelope

containing their allocation and details of procedure in relation to their allocation.

All subjects first underwent a comprehensive interview and full physical examination at the Department of Public Health and Primary Health Care, UiB. This multidimensional examination was important in order to broadly classify each subject based on his or her pain provocative postures and movement behaviours, lifestyle and cognitive behaviours (O'Sullivan, 2005). During the interview, subjects were guided in questioning to inform: their history of pain, pain area and nature, pain behaviour (aggravating/easing movements and activities), their primary functional impairments, disability levels, activity levels and sleep patterns. Inquiries were also made regarding their level of fear of pain and any avoidance of activities, work and social engagement. Their degree of pain focus, pain coping strategies, stress response and its relationship to pain, and their pain beliefs were also questioned as was any history of anxiety and depression. Finally, their beliefs and goals regarding management of their disorder were ascertained. This information was considered alongside the Orebro Musculoskeletal Pain Questionnaire (OMPQ) (Linton and Boersma, 2003) where the patient reported high levels (>5) of stress/anxiety, depressed mood, fear of movement and lack of pain coping strategies. These factors were then considered within the context of the movement behaviours and lifestyle factors, i.e., in the context of fear, avoidance of physical activity and specific pain provocative movement patterns were assessed.

The physical examination involved analysis of the subject's primary functional impairments (pain provocative postures, movements and functional tasks reported during the interview), assessment of their body control and awareness, as well as easing postures and movements (O'Sullivan, 2005).

The physical examination process involved a systematic process of assessment of pain provocative postures (such as sleeping, sitting, standing and bending) and functional movement tasks (such as sit to stand, single-leg stand, spinal movements and lifting) and any other specific tasks nominated by the patient as pain provocative or that they avoided. The validity of this clinical examination approach has been demonstrated in a number of studies (O'Sullivan et al., 2006a,b; Dankaerts et al., 2007; O'Sullivan and Beales, 2007a,b; Beales et al., 2009; Dankaerts et al., 2009; Sheeran et al., 2012). The reliability has previously been reported for both physical and cognitive aspects based on the health-care practitioners' ability to synthesize the patient story and clinical

examination to broadly classify the patient, based on identification of the primary drivers of the disorder within the different levels of the MDCS (Dankaerts et al., 2006; Fersum et al., 2009). The examination process follows a systematic approach (as outlined in Supporting Information Appendix S1), but is tailored to the patients' presentation. In Supporting Information Appendix S2, all the included subjects and their subsequent classification are outlined.

The intervention took place at 3 different private clinics and lasted for 12 weeks. Patients were followed-up immediately and 12 months post-intervention. After the 12-week intervention period, participants were permitted to seek alternative care, and the frequency and type of treatment were monitored in the follow-up questionnaires. A tester blinded to allocation, pretreatment and at 3-month follow-up distributed the questionnaires. During the 12-month follow-up, the questionnaires were sent by mail.

2.3 Interventions

2.3.1 CB-CFT

The behavioural management approach provided was based on the MDCS described by O'Sullivan (2005) and Fersum et al. (2009). Three experienced physiotherapists who conducted the management had undergone, on average, 106 hours of CB-CFT training (including workshops, patient examinations, pilot study and clinical manual). Depending on the different levels of the classification system, each patient received a specific targeted intervention directed at changing their individual cognitive, movement and lifestyle behaviours considered by the therapist to be maladaptive (provocative) of their disorder. Physiotherapists were also guided by the individuals' OMPQ in order to better target their management within the psychosocial domain. The CB-CFT intervention has four main components: (1) a cognitive component, for each patient, their vicious cycle of pain was outlined in a diagram based on their findings from the examination and the OMPQ; (2) specific movement exercises designed to normalize maladaptive movement behaviours as directed by the movement classification; (3) targeted functional integration of activities in their daily life, reported to be avoided or provocative by the patient; and (4) a physical activity programme tailored to the movement classification (see Supporting Information Appendix S3 for detailed description of the CB-CFT). The initial session was 1 h and follow-ups were 30–45 min. Patients were seen at a weekly basis

for the first two to three sessions and then progressed to one session every 2–3 weeks during the 12-week intervention period.

All instructions for subjects were written, and they were asked to complete the programme on a daily basis and complete a daily diary outlining if they had complied with each aspect of the intervention.

2.3.2 MT-EX

Participants allocated to the comparison group were treated with joint mobilization or manipulation techniques applied to the spine or pelvis consistent with best current manual therapy practice. These therapists were specialists in orthopaedic manual therapy with an average of 25.7 years of experience with no prior training in the use of the MDCS or CB-CFT. The particular dose and techniques were at the discretion of the treating therapist, based on each participant's examination findings. In addition, most patients (82.5%) in this group were given exercises or a home exercise programme. This included general exercise or motor control exercise, but not based on the specific MDCS as outlined by O'Sullivan (2005). The motor control exercises involved isolated contractions of the deep abdominal muscles in different functional positions as previously described (Richardson et al., 1998). The therapists in this group generally spent 1 h with the patients for the initial consultation and 30 min for follow-ups.

2.4. Treatment fidelity

Several strategies were applied to enhance treatment fidelity. For every patient, a report was written, documenting type of treatment, number of treatments, medication usage before and after the intervention. In addition, there was a therapist session-by-session documentation of treatment content as well as individual working sheets. The mean number of treatments was 7.7 in the CB-CFT group (range 4–16; SD 2.6), and 8.0 in the MT-EX group (range 3–17; SD 2.9). All the treating physiotherapists in both treatment arms prior to the intervention underwent half a day of training with a clinical psychologist regarding the concepts of best-practice cognitive approach to managing back pain in order to standardize this for both intervention arms as per clinical guidelines. Therapy was in the primary care setting for both groups and was pragmatic. Treatment was discontinued if the therapist deemed the participant had no further need of treatment before the 12 weeks were completed, as is standard clinical practice. There was

no difference between the two groups in terms of medication intake before or after the treatment.

2.5. Outcome measures

2.5.1 ODI

The primary outcome measurement for perceived function was the ODI (Roland and Fairbank, 2000). The ODI psychometric properties have been well established (Fairbank and Pynsent, 2000). Studies have shown good construct validity (Greenough, 1993; Strong et al., 1994), acceptable internal consistency and test-retest reliability (Kopeck et al., 1996), and high responsiveness (Deyo and Centor, 1986).

2.5.2 Pain

Pain intensity in the previous week was another primary outcome, measured by the PINRS (Jensen et al., 1986). The PINRS measures pain severity by asking the patient to select a number (from 0 to 10) to represent how severe the pain had been over the last 2 weeks. PINRS is more reliable than the visual analogue scale, especially with less educated patients (Ferraz et al., 1990).

2.5.3 Hopkins Symptoms Checklist

Secondary outcome measures for anxiety and depression were Hopkins Symptoms Checklist (HSCL-25) (Derogatis et al., 1974). It consists of 25 items: part I of the HSCL-25 has 10 items for anxiety symptoms; part II has 15 items for depression symptoms. Optimal HSCL-25 cut-off was 1.67 for men and 1.75 for women (Sandanger et al., 1998).

2.5.4 Fear-Avoidance Beliefs Questionnaire (FABQ)

The fear-avoidance beliefs of physical activity and work were measured using the FABQ (Waddell et al., 1993). The reliability and validity of this measure has been reported previously (Kovacs et al., 2006).

2.5.5 Total lumbar spine range of motion

Total lumbar spine range of motion was measured with the two inclinometer method (Mayer et al., 1984) utilizing standardized protocol (Adams et al., 1986; Bullock-Saxton, 1993; Chaory et al., 2004). The true lumbar motion was obtained from a subtraction

of the pelvic motion from the gross motion expressed in angular degrees of flexion and extension (Adams et al., 1986).

2.5.6 Patient satisfaction

The patients filled out a patient satisfaction questionnaire (Ware and Davies, 1983) at 3- and 12-month follow-up. This was just a simple questionnaire from 1 to 5 asking the patients how satisfied they were with their treatment: 1 = satisfied, 2 = just a little satisfied, 3 = neither satisfied nor dissatisfied, 4 = just a little dissatisfied, 5 = dissatisfied.

2.5.7 Sick-leave days

In case of sick-leave days, this was extracted from the Ørebro Screening Questionnaire using a 10-category variable. The question asked: 'How many days of work have you missed because of pain during the last 18 months?' The 10 categories were merged into 3. First category was 0 days, second category was 1–7 days and third category was >7 days.

2.5.8 Care-seeking

Questions regarding subsequent treatments between the 3-month intervention period and long-term follow-up were also assessed with a questionnaire at 12 months. The patients were asked the following questions: (1) Have you had any treatment since the intervention finished? Yes/No; (2) What type of treatment have you had?; and (3) How many treatments have you had?

2.6 Statistical considerations and analysis

A linear mixed model was used to estimate the group differences in treatment effect at both time points and also in the change in outcome from 3- and 12-month follow-up, with baseline values included as the only covariate. Age, gender, body mass index (BMI), LBP duration and work status were evaluated as possible confounders, but did not need to be included in the final models. Bootstrapped standard errors were estimated to adjust for departures from normality, as some outcome measures displayed slightly skewed distributions. Models were examined to confirm the absence of influential outlying observations. Statistical significance was set at $p = 0.05$.

Analysis was performed on an 'available case' basis. Two subjects (one CB-CFT and one MT-EX) were missing data at 3-month follow-up but provided




Figure 1 Flow chart depicting participant recruitment and final enrolment for the two groups: manual therapy and exercise (MT-EX) and classification-based cognitive functional therapy (CB-CFT).

12-month data, and five subjects (two CB-CFT and three MT-EX) were missing 12-month follow-up data but provided 3-month data. These cases were included in the model, as the linear mixed model used is a likelihood-based estimation procedure resulting in non-biased estimates, provided data are missing at random. Statistically significant group differences in sick-leave days, patient satisfaction and number of treatments sought post-intervention were assessed using two-sample Wilcoxon rank sum (Mann–Whitney) test. In the case of sick-leave days, the original variable from Ørebro Screening Questionnaire collapsed down for tabular display but analysed using original 10-category variable, i.e., 0, 1–2, 3–7, . . . , etc., days.

In addition, change in pre- to post-treatment was estimated in both treatment groups using paired *t*-tests, and the change in ODI and PINRS were calculated for each study participant, in terms of absolute change from baseline, and tabulated with reference to consensus values for minimally important change (MIC) in these outcomes.

All authors had full access to the data in the study and the corresponding author had final responsibility to submit for publication. An independent person, unfamiliar with the aim, groups and content of the study, entered data into a database. Statistical analysis was performed using Stata/IC 10.1 for Windows (StataCorp LP, College Station, TX, USA). A secondary independent statistician blinded to the treatment

group allocation and unfamiliar with the aim, groups and content of the study also reviewed the statistical analysis.

3. Results

Out of the 169 patients initially enrolled, 121 patients met the inclusion criteria and were found eligible. In the randomized cohort, 62 patients were assigned to the CB-CFT group, and 59 were assigned to the MT-EX group (Fig. 1). A total of 16 of 59 (27.1%) patients assigned to MT-EX and 11 of 62 (17.7%) patients assigned to CB-CFT either did not start treatment or did not complete treatment, and were unavailable for either 3- or 12-month follow-up assessment, which precluded an intention-to-treat analysis. Table 1 displays the baseline characteristics of these patients, by treatment allocation.

Analysed study participants in the two treatment arms were comparable in terms of baseline characteristics, with the exception of small but significant differences in HSCL and FABQ work (Table 1).

Both groups significantly improved with the respective therapeutic interventions. After adjustment for baseline scores, the CB-CFT group displayed superior outcomes supported by both statistically and clinically significant differences when compared with the MT-EX group. This was evident both immediately after and at 12-month post-intervention for both primary and secondary outcomes (Table 2). This was

Table 1 Baseline characteristics of study participants.

Characteristics	Analysed (<i>n</i> = 94)		Excluded from analysis (<i>n</i> = 27) (did not start or discontinued treatment)	
	MT-EX (<i>n</i> = 43)	CB-CFT (<i>n</i> = 51)	MT-EX (<i>n</i> = 16)	CB-CFT (<i>n</i> = 11)
Age – mean (SD)	42.9 (12.5)	41.0 (10.3)	39.9 (13.8)	37.0 (10.5)
Female – <i>n</i> (%)	21 (48.8)	27 (52.9)	10 (62.5)	5 (45.5)
LBP duration – <i>n</i> (%) ^a				
3–12 months	6 (14.3)	6 (11.8)	0	0
1–5 years	13 (31.0)	14 (27.5)	4 (30.8)	1 (12.5)
>5 years	23 (54.8)	31 (60.8)	9 (69.2)	7 (87.5)
Height (cm) (SD)	173.5 (8.6)	174.3 (9.4)	171.8 (10.1)	177.4 (7.9)
Weight (SD)	76.2 (13.1)	78.3 (16.6)	76.9 (16.5)	85.4 (11.9)
BMI (SD)	25.2 (3.5)	25.6 (4.0)	25.9 (4.3)	27.1 (3.1)
Work status				
Paid work <i>n</i> (%)	34 (79.1)	45 (88.2)	15 (93.8)	11 (100.0)
Primary outcomes – mean (SD)				
Oswestry Disability Index	24.0 (8.0)	21.3 (7.5)	27.9 (7.6)	23.1 (8.1)
Pain intensity last week	5.3 (1.9)	4.9 (2.0)	6.2 (1.2)	6.0 (1.9)
Secondary outcomes – mean (SD)				
Hopkins Symptoms Checklist	1.57 (0.39)	1.40 (0.33)	1.78 (0.53)	1.60 (0.42)
Fear-avoidance physical	11.8 (5.0)	11.1 (3.9)	12.6 (4.6)	10.9 (5.3)
Fear-avoidance work	19.3 (11.1)	14.1 (9.6)	17.8 (8.2)	13.5 8.7
Sick-leave days – <i>n</i> (%)				
None	13 (30.2)	15 (29.4)	2 (12.5)	3 (27.3)
1–7 days	9 (20.9)	13 (25.5)	4 (25.0)	2 (18.2)
>7 days	21 (48.9)	23 (45.1)	10 (62.5)	6 (54.5)
Lumbar ROM – mean (SD)	46.2 (13.0)	50.2 (14.9)	44.3 (13.6)	57.6 (9.8)

Disability: Oswestry Disability Index (0–100), low scores indicate low disability.

Pain: intensity last week (pain intensity numerical rating scale 0–10), low scores indicate low pain intensity.

Anxiety/depression: Hopkins Symptoms Checklist – 25-item checklist – low scores indicate low anxiety and depression.

Fear of movement/(re) injury: The Fear-Avoidance Beliefs – Physical (0–24), low scores indicate low fear related to physical activities.

Fear of movement/(re) injury: The Fear-Avoidance Beliefs – Work (0–42), low scores indicate low fear related to work activities.

Range of motion: Lumbar motion was obtained from a subtraction of the pelvic motion from the gross motion expressed in angular degrees of flexion and extension.

BMI, body mass index; CB-CFT, classification-based cognitive functional therapy; LBP, low back pain; MT-EX, manual therapy and exercise; ROM, range of motion; SD, standard deviation.

^aMissing four cases from MT-EX and three from CB-CFT group.

demonstrated by the degree of improvement in the CB-CFT group for ODI score being 13.7 points [95% confidence interval (CI): 11.4–16.1; $p < 0.001$] and for PINRS scores 3.2 (95% CI: 2.5–3.9; $p < 0.001$). In the MT-EX group, the mean improvement for ODI score was 5.5 points (95% CI: 2.8–8.3; $p < 0.001$) and 1.5 for PINRS (95% CI: 0.7–2.2; $p < 0.001$). The improvements for all secondary outcomes showed similar effects, with the CB-CFT group demonstrating significantly greater change when compared with the MT-EX group across all outcomes, except for total lumbar range of motion (Table 2).

Table 3 presents the improvements in primary outcome measures in terms of absolute improvement from baseline. There was maintenance of treatment effect over the 3- to 12-month follow-up time for both

groups, with no significant main effect for time or group/time interaction identified in the linear mixed models.

4. Discussion

This study reveals that the CB-CFT group demonstrated superior outcomes compared with the MT-EX group across every domain measured at post-intervention and at 12-month follow-up (Table 2). Both groups showed significant improvement in short- and long-term follow-ups; however, the CB-CFT group was superior based on clinically meaningful changes as defined by MIC, defined as >10-point change in ODI and >1.5 on PINRS (Ostelo et al., 2008).

Table 2 Outcomes (unadjusted means and SDs).

	MT-EX		CB-CFT		CB-CFT vs MT-EX ^a Mean difference (95 % CI)						
	Mean	SD	Mean	SD							
Primary outcome variables											
Oswestry Disability Index Questionnaire											
Baseline	24.0	8.0	21.3	7.5							
3 months	18.5	8.1	7.6	6.7	-9.7 (-12.7 to -6.7)***						
12 months	19.7	11.7	9.9	9.8	-8.2 (-12.6 to -3.8)***						
Pain intensity in last week											
Baseline	5.3	1.9	4.9	2.0							
3 months	3.8	1.9	1.7	1.7	-2.1 (-2.7 to -1.4)***						
12 months	3.8	2.1	2.3	2.0	-1.3 (-2.1 to -0.5)***						
Secondary outcome variables											
Hopkins Symptoms Checklist											
Baseline	1.56	0.39	1.40	0.33							
3 months	1.43	0.37	1.20	0.27	-0.12 (-0.19 to -0.04)**						
12 months	1.51	0.47	1.22	0.32	-0.13 (-0.22 to -0.04)**						
Fear-avoidance physical											
Baseline	11.8	5.0	11.1	3.9							
3 months	10.3	6.0	6.1	5.0	-3.6 (-5.3 to -1.9)***						
12 months	10.9	5.5	5.8	5.5	-4.7 (-6.5 to -3.0)***						
Fear-avoidance work											
Baseline	19.1	11.1	14.1	9.6							
3 months	17.4	10.8	8.3	8.4	-5.7 (-7.8 to -3.6)***						
12 months	16.6	12.2	7.7	9.0	-5.6 (-8.7 to -2.5)***						
Total lumbar spine range of motion											
Baseline (degrees)	46.2	13.0	50.2	14.9							
3 months	45.6	12.7	49.7	14.0	1.9 (-2.8 to 6.7)						
Sick-leave days [n (%)]											
	MT-EX			CB-CFT							
	0	1-7	>7	0	1-7	>7					
Baseline	14 (31.8)	9 (20.5)	21 (47.7)	15 (29.4)	13 (25.5)	23 (45.1)					
12 months	16 (40.0)	7 (17.5)	17 (42.5)	32 (65.3)	7 (14.3)	10 (20.4)	z = 2.95**				
Patient satisfaction^b [n (%)]											
	MT-EX					CB-CFT					
	1	2	3	4	5	1	2	3	4	5	
3 months ³	28 (68.3)	6 (14.6)	5 (12.2)	0	2 (5.9)	48 (94.1)	1 (2.0)	2 (3.9)	0	0	z = 3.21**
12 months ³	18 (46.2)	8 (20.5)	9 (23.1)	3 (7.7)	1 (2.5)	46 (95.8)	1 (2.1)	1 (2.1)	0	0	z = 5.18***
Care-seeking after intervention [n (SD)]											
	MT-EX					CB-CFT					
12 months	10.6 (13.3)					2.1 (5.4)					z = 4.79***

Disability: Oswestry Disability Index (0–100), low scores indicate low disability.

Pain: intensity last week (pain intensity numerical rating scale 0–10), low scores indicate low pain intensity.

Anxiety/depression: Hopkins Symptoms Checklist – 25-item checklist – low scores indicate low anxiety and depression.

Fear of movement/(re) injury: The Fear-Avoidance Beliefs – Physical (0–24), low scores indicate low fear related to physical activities.

Fear of movement/(re) injury: The Fear-Avoidance Beliefs – Work (0–42), low scores indicate low fear related to work activities.

Range of motion: Lumbar motion was obtained from a subtraction of the pelvic motion from the gross motion expressed in angular degrees of flexion and extension.

CB-CFT, classification-based cognitive functional therapy; MT-EX, manual therapy and exercise; SD, standard deviation.

^aNegative values favour CB-CFT.

^bPatient satisfaction: (1–5), 1 = completely satisfied, 2 = a little bit satisfied, 3 = neither satisfied or dissatisfied, 4 = a little bit dissatisfied, 5 = completely dissatisfied.

**p < 0.01.

***p < 0.001.

Table 3 Immediate and long-term improvements in primary outcome measures from baseline [*n* (%)].

	3 months				12 months			
	MT-EX		CB-CFT		MT-EX		CB-CFT	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Oswestry								
No change	14	33.3	2	4.0	12	30.8	5	10.2
<10 points	14	33.3	12	24.0	15	38.5	13	26.5
10–19 points	12	28.6	23	46.0	9	23.1	24	49.0
20–29 points	1	2.4	11	22.0	2	5.1	5	10.2
30 or more points	1	2.4	2	4.0	1	2.5	2	4.1
Pain (PINRS)								
≤0	15	35.7	6	12.0	12	30.8	10	20.4
1 point	8	19.1	6	12.0	11	28.2	3	6.1
2 points	7	16.7	9	18.0	7	17.9	14	28.6
3 points	4	9.5	8	16.0	4	10.3	7	14.3
4–5 points	4	9.5	13	26.0	3	7.7	7	14.3
>6 points	4	9.5	8	16.0	2	5.1	8	16.3

Minimally important change: Clinically meaningful changes are defined as >10-point change in Oswestry Disability Index and >1.5 on PINRS (Ostelo et al., 2008). CB-CFT, classification-based cognitive functional therapy; MT-EX, manual therapy and exercise; PINRS, pain intensity numerical rating scale.

The effect sizes of conservative interventions from previous Cochrane reviews reveal findings similar to the MT-EX group (Assendelft et al., 2004). In response, calls have been made for a paradigm shift, away from a biomedical ‘injury’ model, to viewing LBP as a multifactorial biopsychosocial disorder, and directing treatment at beliefs and behaviours that promote pain and disability rather than simply at the signs and symptoms associated with the disorder. Calls have also been made for the need for a multidimensional classification-based approach to direct management of NSCLBP in order to make treatment more person-centred (Borkan et al., 2002; O’Sullivan, 2011). This is supported by reports that disability levels in chronic pain are better predicted by cognitive and behavioural aspects of pain rather than sensory and biomedical ones (Campbell and Edwards, 2009). CB-CFT addresses all of these objectives.

Although satisfaction rates were high in both groups, odds of being completely satisfied were over three times higher in the CB-CFT group at 3 months and five times higher at 12 months. The degree of patient satisfaction is seen as a reflection of the quality of care and as an important outcome in its own right (May, 2001). The importance of communication, patient-centred approaches and goal matching has been well documented in the literature as important for the therapeutic relationship, enhancing compliance and patient outcomes (Linton, 2005; O’Sullivan, 2011).

CB-CFT had a strong cognitive focus with an emphasis on reframing the persons’ understanding of their back pain in a person-centred manner, with an emphasis on changing maladaptive movement, cognitive and lifestyle behaviours contributing to their vicious cycle of pain. This was performed by means of reflective communication, providing a contemporary understanding of pain mechanisms, correcting faulty pathoanatomical beliefs, goal setting, verbal, written and visual feedback (viewing their own back) and a strong emphasis on normalizing movement behaviours within a graded functional approach.

Although this is the first RCT to address maladaptive movement behaviours specific to the patient’s presentation within a cognitive framework, the exact benefits from targeting specific movement training cannot be isolated from the other aspects of the intervention. The behaviours that were targeted were prioritized based on the movements or postures that patients reported that they most feared, avoided and/or that provoked them. These identified movements were the targets for the movement retraining aspect of the intervention based on the patient’s classification and were integrated to the goals of the patient. The use of visual feedback such as mirrors was central to this process. The aim of this approach was for each subject to acquire self-management strategies for their disorder by developing positive back pain beliefs, pain control, developing adaptive strategies of movement that enhanced functional capacity and the ability to engage in regular physical activity.

These findings are supported by previous reports of benefits with different targeted behavioural approaches to managing NSCLBP. Moseley et al. (2004) reported reduced pain and enhanced function associated with pain education. Asenlof et al. (2009) reported superior long-term outcomes for treating NSCLBP with an individually tailored behavioural intervention targeting cognitions, motor behaviour and activity, compared with physical therapy. The use of visual feedback when training movement in patients with LBP has also been shown to reduce pain and influence functional capacity (Sheeran et al., 2012; Wand et al., 2012). Hill et al. (2011) reported superior outcomes when management was targeted on the basis of psychosocial risk factors. Vlaeyen et al. (2002) reported benefits with a graded exposure approach to management in a series of NSCLBP patients with high levels of fear. CB-CFT incorporates all of these aspects within its intervention.

Given the multidimensional nature of the CB-CFT intervention, it is not clear as to the exact basis for the superior outcomes. We hypothesize that the

mechanisms for change are likely to be multifactorial given the patient-centred body–mind behavioural approach, in contrast to a more treatment-orientated signs and symptoms approach in manual therapy. On one hand, this behavioural approach may have impacted on cognitive factors known to affect pain sensitivity and disability such as developing positive beliefs, reduced fear, increased awareness, enhanced understanding and control of pain, adaptive coping, enhanced self-efficacy, confidence and improved mood. Evidence for this is supported by the reduction in fear of movement and improved mood observed following the intervention. On the other hand, the functional behavioural aspects of the intervention were targeted at enhancing body awareness, relaxation of guarded muscles, normalizing maladaptive movement patterns, body schema retraining with the use of mirror feedback, extinguishing pain behaviours, conditioning and increased functional capacity. These factors have been associated with levels of pain, disability, fear and catastrophizing (Wideman et al., 2009; Lewis et al., 2012; Wand et al., 2012). We also acknowledge that the active engagement required of subjects for this behavioural approach may present a barrier for those unwilling to self-manage their disorder (Carr et al., 2006). This may be dependent on the levels of acceptance and readiness to engage in behavioural change, although these factors were not formally assessed.

Although it was not a primary aim of the CB-CFT intervention, the results demonstrate a 2.95-times less likelihood of taking sick leave for their LBP at 12 months compared with the MT-EX group. Previously, only studies using cognitive behavioural therapy in multidisciplinary treatment models have shown an effect on sick-listing for this patient group (Airaksinen et al., 2006). However, as these numbers were extracted from the OMPQ and were self-reported, these findings must be interpreted with some caution. The patients in the CB-CFT group also sought less additional treatment for their pain, implying they may have been more empowered to self-manage their disorder, suggesting significant cost–benefits.

A limitation of this study was the number of patients that either did not start or complete treatment. While there was a comparable proportion of non-completers in each group (Table 2), 8 of 59 (13.5%) of the MT-EX group failed to commence their allocated treatment, compared with only 1 of 62 (1.6%) of the CB-CFT group. This may be due to the fact that seven of these subjects had reported previous manual therapy treatment with poor effect, which would have potentially biased for a poorer outcome in

the MT-EX group. It should be noted that there was no statistically significant difference between completers and non-completer based on baseline characteristics, and as we performed our analysis conditioning on baseline scores and confirmed the absence of a confounding effect of age, gender, BMI, LBP duration and work status on our results, we are confident that our estimates of treatment effect are not substantially biased by these missing cases. Also of note, no drop-outs reported adverse effects from either intervention arm. Furthermore, due to a lack of power, we were unable to determine the influence of the subject classification on the outcome.

We acknowledge that there are also a number of additional methodological considerations that may have influenced the results of this study. Firstly, the patients were recruited from a variety of sources both primary and secondary care levels, as well as newspaper advertising, which could have influenced the kind of patients who entered the study. However, the wide inclusion criteria in the study suggest that a common and representative group of patients with chronic localized LBP without objective sign of pathology to the spine were included. Webb et al. (2003) reported that an Oswestry score of >25% is considered the cut-off score for classifying ‘disabling back pain’. The patients recruited were, on average, just below this [24.0 (MT-EX) and 21.3 (CB-CFT)] at baseline, and hence it can be said that the patients sampled in this study had moderate back pain and functional impairment sufficient to result in activity limitation and sick leave for many (see Table 1). It is also acknowledged that therapists in both arms of the study were not blinded to the intervention, and although all therapists had considerable experience, the influence of therapist enthusiasm and expectation for change was not controlled for.

Furthermore, the multidimensional nature of the study limits any conclusion as to the specific effects of the different components of the intervention. Future research that investigates matching versus non-matching of interventions for patients with chronic mechanical LBP may help identify the effects of specific aspects of the intervention (Kent et al., 2010). Also given this was a pragmatic trial, the intervention dose was not controlled in either group, although both groups received remarkably similar attention. While this intervention appears to be successful for the population we tested, further studies are needed to investigate those with higher levels of pain and disability, patients that are long-term sick-listed as well as in other cultural and occupational groups, in order to determine the generalizability of the findings. This

approach also needs to be compared with other cognitive approaches and tested within a multi-centre trial framework in the future.

4.1 Conclusions

The results of this study support that a behaviourally orientated targeted approach to manage NSCLBP (CB-CFT) was more effective at reducing pain, disability, fear beliefs, mood and sick leave at long-term follow-up than MT-EX.

Acknowledgements

Research fellow Fersum was supported by the Norwegian Fund for Post-Graduate Training in Physiotherapy.

Many thanks to the treating therapist in both treatment arms AB, RH, NE, EØ and LGL.

Author contributions

K.V.F., P.O., J.S.S. and A.K. were all involved in the planning of the RCT, preparing protocols, ethical approval and deciding the interventions utilized. A.S. was involved in the statistical methods used in the manuscript. K.V.F., P.O., J.S.S., A.S. and A.K. were all involved in the writing and editing the manuscript. All authors discussed the results and commented on the manuscript.

References

- Adams, M., Dolan, P., Marx, C., Hutton, W. (1986). An electronic inclinometer technique for measuring lumbar curvature. *Clin Biomech* 1, 130–134.
- Airaksinen, O., Brox, J.I., Cedraschi, C., Hildebrandt, J., Klaber-Moffett, J., Kovacs, F., Mannion, A.F., Reis, S., Staal, J.B., Ursin, H., Zanoli, G. (2006). Chapter 4. European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J* 15(Suppl 2), S192–S300.
- Asenlof, P., Denison, E., Lindberg, P. (2005). Individually tailored treatment targeting activity, motor behavior, and cognition reduces pain-related disability: A randomized controlled trial in patients with musculoskeletal pain. *J Pain* 6, 588–603.
- Asenlof, P., Denison, E., Lindberg, P. (2009). Long-term follow-up of tailored behavioural treatment and exercise based physical therapy in persistent musculoskeletal pain: A randomized controlled trial in primary care. *Eur J Pain* 13, 1080–1088.
- Assendelft, W.J.J., Morton, S.J., Yu, E.I., Suttorp, M.J., Shekelle, P.G. (2004). Spinal manipulative therapy for low back pain. *Cochrane Database Syst Rev* (1)CD000447.
- Beales, D.J., O'Sullivan, P.B., Briffa, N.K. (2009). Motor control patterns during an active straight leg raise in chronic pelvic girdle pain subjects. *Spine* 34, 861–870.
- Bjorck-van Dijken, C., Fjellman-Wiklund, A., Hildingsson, C. (2008). Low back pain, lifestyle factors and physical activity: A population-based study. *J Rehabil Med* 40, 864–869.
- Boersma, K., Linton, S.J. (2006). Psychological processes underlying the development of a chronic pain problem: A prospective study of the relationship between profiles of psychological variables in the fear-avoidance model and disability. *Clin J Pain* 22, 160–166.
- Borkan, J., Van Tulder, M., Reis, S., Schoene, M.L., Croft, P., Hermoni, D. (2002). Advances in the field of low back pain in primary care: A report from the fourth international forum. *Spine* 27, E128–E132.
- Bouter, L., Penick, V., Bombardier, C. (2003). Cochrane back review group. *Spine* 28, 1215–1218.
- Bullock-Saxton, J. (1993). Postural alignment: A repeatability study. *Aust J Physiother* 39, 25–29.
- Campell, C., Edwards, R. (2009). Mind-body interactions in pain: The neurophysiology of anxious and catastrophic pain-related thoughts. *Transl Res* 153, 97–101.
- Carr, J.L., Moffett, J.A., Sharp, D.M., Haines, D.R. (2006). Is the Pain Stages of Change Questionnaire (PSOCQ) a useful tool for predicting participation in a self-management programme? Further evidence of validity, on a sample of UK pain clinic patients. *BMC Musculoskelet Disord* 7, 101.
- Chaory, K., Fayad, F., Rannou, F., Lefevre-Colau, M.M., Fermanian, J., Revel, M., Poiraudou, S. (2004). Validation of the French version of the fear avoidance belief questionnaire. *Spine* 29, 908–913.
- Dankaerts, W., O'Sullivan, P.B., Burnett, A.F., Straker, L.M. (2007). The use of a mechanism-based classification system to evaluate and direct management of a patient with non-specific chronic low back pain and motor control impairment – a case report. *Man Ther* 12, 181–191.
- Dankaerts, W., O'Sullivan, P.B., Burnett, A.F., Straker, L.M., Davey, P., Gupta, R. (2009). Discriminating healthy controls and two clinical subgroups of nonspecific chronic low back pain patients using trunk muscle activation and lumbosacral kinematics of postures and movements: A statistical classification model. *Spine* 34, 1610–1618.
- Dankaerts, W., O'Sullivan, P.B., Straker, L.M., Burnett, A.F., Skouen, J.S. (2006). The inter-examiner reliability of a classification method for non-specific chronic low back pain patients with motor control impairment. *Man Ther* 11, 28–39.
- Derogatis, L., Lipman, R., Rickels, K., Uhlenhuth, E., Covi, L. (1974). The Hopkins Symptom Checklist (HSCL): A self-report symptom inventory. *Behav Sci* 19, 1–15.
- Deyo, R.A., Centor, R.M. (1986). Assessing the responsiveness of functional scales to clinical change: An analogy to diagnostic test performance. *J Chronic Dis* 39, 897–906.
- Fairbank, J.C.T., Pynsent, P.B. (2000). The Oswestry Disability Index. *Spine* 25, 2940–2953.
- Ferraz, M.B., Quaresma, M.R., Aquino, L.R., Atra, E., Tugwell, P., Goldsmith, C.H. (1990). Reliability of pain scales in the assessment of literate and illiterate patients with rheumatoid arthritis. *J Rheumatol* 17, 1022–1024.
- Fersum, K.V., Dankaerts, W., O'Sullivan, P.B., Maes, J., Skouen, J.S., Bjordal, J.M., Kvale, A. (2010). Integration of subclassification strategies in randomised controlled clinical trials evaluating manual therapy treatment and exercise therapy for non-specific chronic low back pain: A systematic review. *Br J Sports Med* 44, 1054–1062.
- Fersum, K.V., O'Sullivan, P.B., Kvale, A., Skouen, J.S. (2009). Inter-examiner reliability of a classification system for patients with non-specific low back pain. *Man Ther* 14, 555–561.
- Furlan, A.D., van Tulder, M.W., Cherkin, D.C., Tsukayama, H., Lao, L., Koes, B.W., Berman, B.M. (2005). Acupuncture and

- dry needling for low back pain. *Cochrane Database Syst Rev* (1)CD001351.
- Greenough, C.G. (1993). Recovery from low back pain. 1–5 year follow-up of 287 injury-related cases. *Acta Orthop Scand Suppl* 254, 1–34.
- Hayden, J.A., van Tulder, M.W., Tomlinson, G. (2005). Systematic review: Strategies for using exercise therapy to improve outcomes in chronic low back pain. *Ann Intern Med* 142, 776–785.
- Hill, J.C., Dunn, K.M., Main, C.J., Hay, E.M. (2010). Subgrouping low back pain: A comparison of the STarT Back Tool with the Orebro Musculoskeletal Pain Screening Questionnaire. *Eur J Pain* 14, 83–89.
- Hill, J.C., Whitehurst, D.G., Lewis, M., Bryan, S., Dunn, K.M., Foster, N.E., Konstantinou, K., Main, C.J., Mason, E., Somerville, S., Sowden, G., Vohora, K., Hay, E.M. (2011). Comparison of stratified primary care management for low back pain with current best practice (STarT Back): A randomised controlled trial. *Lancet* 378, 1560–1571.
- Jensen, M.P., Karoly, P., Braver, S. (1986). The measurement of clinical pain intensity: A comparison of six methods. *Pain* 27, 117–126.
- Kent, P., Hancock, M., Petersen, D.H., Mjosund, H.L. (2010). Clinimetrics corner: Choosing appropriate study designs for particular questions about treatment subgroups. *J Man Manip Ther* 18, 147–152.
- Klaber Moffett, J.A., Carr, J., Howarth, E. (2004). High fear-avoiders of physical activity benefit from an exercise program for patients with back pain. *Spine* 29, 1167–1172.
- Kopec, J.A., Esdaile, J.M., Abrahamowicz, M., Abenhaim, L., Wood-Dauphinee, S., Lamping, D.L., Williams, J.I. (1996). The Quebec Back Pain Disability Scale: Conceptualization and development. *J Clin Epidemiol* 49, 151–161.
- Kovacs, F.M., Muriel, A., Medina, J.M., Abaira, V., Sanchez, M.D., Jauregui, J.O. (2006). Psychometric characteristics of the Spanish version of the FAB questionnaire. *Spine* 31, 104–110.
- Lewis, S., Holmes, P., Woby, S., Hindle, J., Fowler, N. (2012). The relationships between measures of stature recovery, muscle activity and psychological factors in patients with chronic low back pain. *Man Ther* 17, 27–33.
- Linton, S.J. (2000). A review of psychological risk factors in back and neck pain. *Spine* 25, 1148–1156.
- Linton, S.J. (2005). *Understanding Pain for Better Clinical Practice: A Psychological Perspective (Pain Research and Clinical Management)*, 1st edn (Edinburgh: Elsevier).
- Linton, S.J., Boersma, K. (2003). Early identification of patients at risk of developing a persistent back problem: The predictive validity of the Orebro Musculoskeletal Pain Questionnaire. *Clin J Pain* 19, 80–86.
- May, S. (2001). Patient satisfaction with management of back pain. *Physiotherapy* 87, 4–20.
- Mayer, T.G., Tencer, A.F., Kristoferson, S., Mooney, V. (1984). Use of non-invasive techniques for quantification of spinal range-of-motion in normal subjects and chronic low-back dysfunction patients. *Spine* 9, 588–595.
- Mitchell, T., O’Sullivan, P., Burnett, A., Straker, L., Smith, A., Thornton, J., Rudd, C.J. (2010). Identification of modifiable personal factors that predict new-onset low back pain: A prospective study of female nursing students. *Clin J Pain* 26, 275–283.
- Moseley, G.L., Nicholas, M.K., Hodges, P.W. (2004). A randomized controlled trial of intensive neurophysiology education in chronic low back pain. *Clin J Pain* 20, 324–330.
- O’Sullivan, P. (2005). Diagnosis and classification of chronic low back pain disorders: Maladaptive movement and motor control impairments as underlying mechanism. *Man Ther* 10, 242–255.
- O’Sullivan, P. (2011). It’s time for change with the management of non-specific chronic low back pain. *Br J Sports Med* 46, 224–227.
- O’Sullivan, P.B., Beales, D.J. (2007a). Diagnosis and classification of pelvic girdle pain disorders – part 1: A mechanism based approach within a biopsychosocial framework. *Man Ther* 12, 86–97.
- O’Sullivan, P.B., Beales, D.J. (2007b). Diagnosis and classification of pelvic girdle pain disorders, part 2: Illustration of the utility of a classification system via case studies. *Man Ther* 12, e1–e12.
- O’Sullivan, P.B., Dankaerts, W., Burnett, A.F., Booth, R., Carlsen, C., Chen, D., Schultz, A. (2006a). Evaluation of the flexion relaxation phenomenon in the trunk muscles in sitting. *Spine* 31, 2009–2016.
- O’Sullivan, P.B., Mitchell, T., Bulich, P., Waller, R., Holte, J. (2006b). The relationship between posture and back muscle endurance in industrial workers with flexion-related low back pain. *Man Ther* 11, 264–271.
- Ostelo, R.W., Deyo, R.A., Stratford, P., Waddell, G., Croft, P., Von Korf, M., Bouter, L., De Vet, H.C. (2008). Interpreting change scores for pain and functional status in low back pain: Towards international consensus regarding minimal important change. *Spine* 33, 90–94.
- Ostelo, R.W., Van Tulder, M.W., Vlaeyen, J.W., Linton, S.J., Morley, S.J., Assendelft, W.J. (2005). Behavioural treatment for chronic low back pain. *Cochrane Database Syst Rev* (1)CD002014.
- Richardson, C., Jull, G., Hodges, P., Hides, J. (1998). *Therapeutic Exercise for Spinal Segmental Stabilization in Low Back Pain. Scientific Basis and Clinical Approach* (London: Churchill Livingstone).
- Riipinen, M., Niemisto, L., Lindgren, K.A., Hurri, H. (2005). Psychosocial differences as predictors for recovery from chronic low back pain following manipulation, stabilizing exercises and physician consultation or physician consultation alone. *J Rehabil Med* 37, 152–158.
- Roland, M., Fairbank, J. (2000). The Roland-Morris Disability Questionnaire and the Oswestry Disability Questionnaire. *Spine* 25, 3115–3124.
- Sandanger, I., Moum, T., Ingebrigtsen, G., Dalgard, O.S., Sorensen, T., Bruusgaard, D. (1998). Concordance between symptom screening and diagnostic procedure: The Hopkins Symptom Checklist-25 and the Composite International Diagnostic Interview I. *Soc Psychiatry Psychiatr Epidemiol* 33, 345–354.
- Sheeran, L., Sparkes, V., Catterson, B., Busse-Morris, M., van Deursen, R. (2012). Spinal position sense and trunk muscle activity during sitting and standing in nonspecific chronic low back pain: Classification analysis. *Spine* 37, E486–E495.
- Smart, K.M., Blake, C., Staines, A., Doody, C. (2011). The discriminative validity of ‘nociceptive,’ ‘peripheral neuropathic,’ and ‘central sensitisation’ as mechanisms-based classifications of musculoskeletal pain. *Clin J Pain* 27, 655–663.
- Spitzer, W. (1987). Scientific approach to the assessment and management of activity related spinal disorders. *Spine* 7, S1–S55.
- Staal, J.B., de Bie, R., de Vet, H.C.W., Hildebrandt, J., Nelemans, P. (2008). Injection therapy for subacute and chronic low-back pain. *Cochrane Database Syst Rev* (3)CD001824.

- Strong, J., Ashton, R., Large, R.G. (1994). Function and the patient with chronic low back pain. *Clin J Pain* 10, 191–196.
- Turk, D.C. (2005). The potential of treatment matching for subgroups of patients with chronic pain: Lumping versus splitting. *Clin J Pain* 21, 44–55.
- Vlaeyen, J.W., Crombez, G. (1999). Fear of movement/(re)injury, avoidance and pain disability in chronic low back pain patients. *Man Ther* 4, 187–195.
- Vlaeyen, J.W., De Jong, J.R., Onghena, P., Kerckhoffs-Hanssen, M., Kole-Snijders, A.M. (2002). Can pain-related fear be reduced? The application of cognitive-behavioural exposure in vivo. *Pain Res Manag* 7, 144–153.
- Vollenbroek-Hutten, M.M.R., Hermens, H.J., Wever, D., Gorter, M., Rinket, J., Ijzerman, M. (2004). Differences in outcome of a multidisciplinary treatment between subgroups of chronic low back pain patients defined using two multiaxial assessment instruments: The Multidimensional Pain Inventory and lumbar dynamometry. *Clin Rehabil* 18, 556–569.
- Waddell, G., Newton, M., Henderson, I., Somerville, D., Main, C.J. (1993). A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain* 52, 157–168.
- Wand, B.M., Tulloch, V.M., George, P.J., Smith, A.J., Goucke, R., O'Connell, N.E., Moseley, G.L. (2012). Seeing it helps: Movement-related back pain is reduced by visualization of the back during movement. *Clin J Pain* 28, 602–608.
- Ware, J., Davies, A. (1983). Defining and measuring patient satisfaction with medical care. *Eval Program Plann* 6, 247–263.
- Webb, R., Brammah, T., Lunt, M., Urwin, M., Allison, T., Symmons, D. (2003). Prevalence and predictors of intense, chronic, and disabling neck and back pain in the UK general population. *Spine* 28, 1195–1202.
- Wideman, T.H., Adams, H., Sullivan, M.J.L. (2009). A prospective sequential analysis of the fear-avoidance model of pain. *Pain* 145, 45–51.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix S1. Displays an overview of the multidimensional classification system. This system is guided by a clinical reasoning process that considers the presence and dominance of different factors, allowing for patient-centred targeted intervention.

Appendix S2. Displays an overview of the randomized subjects ($n = 121$) and how they spread out in the different categories.

Appendix S3. Schematic overview of the four different components of the CB-CFT treatment.

Component	Description All patients were asked to fill in a compliance questionnaire regarding the four aspects of the intervention and present it each session.
1. Cognitive component	<p>For each patient their vicious cycle of pain was outlined in a diagram based on their findings from the examination and the Orebro MPQ. For patients identified with more contributing psychosocial factors, greater emphasis was placed on this aspect of the intervention.</p> <p>For example, it was explained how negative beliefs about pain, fear of movement, increased focus on pain, low mood and poor pacing reinforced avoidance and protective behaviours, muscle guarding, altered movement patterns and body postures, which in turn fed a vicious cycle of pain sensitisation and disability. This process was openly discussed and the patient was invited to consider how they might be able to break their cycle and set their goals for management. These functional goals formed the basis on which the individual exercise management was developed and targeted in the context of lifestyle factors relevant to the individual. Patients were also informed regarding neuro-physiological concepts of pain sensitisation, and that pain does not equal harm or structural damage. If patients had concerns regarding their radiological imaging, they were informed that radiological findings are common in pain free subjects, and correlate poorly with levels of pain and disability.</p> <p>This aspect of the intervention was revisited throughout the treatment periods and integrated into the functional aspects of management as the patient was challenged and exposed to previously provocative tasks. For example with fear of movement, the patients beliefs about pain, perceived consequences and their behavioural responses to both the thoughts and act of the movement were examined. This allowed the patient to be mindful of fearful thoughts driving stress responses and protective behaviours during specific postures, movements and tasks. This reflection allowed them to modify these responses. All other aspects of the intervention listed below had a strong cognitive focus with an emphasis on reflective communication, self-management practices, functional enhancement and goal orientation.</p> <p>Prior to discharge acute exacerbation management planning was discussed with each patient in order to promote an active / confrontation approach to pain management.</p>
2. Functional movement exercises	<p>The aim of this aspect of the intervention was to provide patients with alternative strategies to normalise their postural and movement behaviours.</p> <p>If a patient was unable to relax their trunk muscles, they were initially instructed to learn diaphragm breathing in relaxed postures such as lying, sitting and standing as a first step before movement training.</p> <p>Once this was achieved, all patients received targeted functional postural and movement training based on their movement classification and directed by the activities and postures that they either avoided due to pain or that provoked their pain.</p> <p>This approach followed a ‘graded exposure’ model where the patient was exposed to previously pain provocative tasks, but in a relaxed and controlled manner, as dictated by their movement classification and with feedback (visual with the use of mirrors, mental imagery and awareness of body responses such as breath holding, muscle guarding and changes respiration rate).</p> <p>For example, for those with a ‘movement impairment’ classification, patients were directed to move into the direction of pain provocation in a graded manner (first using less threatening movements and progressing to more threatening movements and activities) in order to restore normal unguarded movement. In the case of flexion ‘movement impairment’ disorder, this first involved gentle non-weight bearing lumbar flexion exercises progressing to flexion in sitting, standing and lifting, whilst ensuring that the movements were performed in a relaxed manner,</p>

without breath holding and protective behaviours. For those with a **‘control impairment’** classification, subjects were trained to modify their pain provocative postures and movement patterns to reduce pain whilst performing the task. For example, for a flexion ‘control impairment’ provoked with sitting, bending and lifting, patients were taught to change their pattern of movement to reduce lumbo-sacral flexion during these tasks. They were first taught to dissociate lumbo-pelvic from thoracic movement in sitting to reduce lumbo-pelvic flexion. This was then progressed to bending and lifting with a focus on facilitating hip/pelvic and thoracic flexion in a relaxed manner (O’Sullivan, 2005a). The management of the few subjects with pelvic girdle pain, followed a similar approach, documented previously (O’Sullivan and Beales, 2007a, b).

Simple non-threatening low load exercises were gradually progressed towards higher load and more complex functional exercises, as the subject gained confidence and control in performing the tasks. No more than 3 or 4 exercises were given at a time.

This challenged each patient to perform functional activities and postures that they nominated as pain and/or fear provoking without adopting pain behaviours such as grimacing, breath holding, muscle guarding, propping with hands or avoidance such as asymmetrical loading.

In this manner they were instructed to change old pain provocative movement behaviours and to reinforce their new functionally enhancing movement behaviours. This was augmented by dynamic practical demonstration of the postures or movement by the therapist, the use of mirrors so they could view their own spines to enhance body schema awareness, written instruction and stick body diagrams (outlining the ‘old way’ vs. the ‘new way’ of sitting, standing, bending, lifting and moving).

Pain related movement behaviours such as propping on the hands, breath holding and abdominal bracing were discouraged.

Where the patient’s functional goals required it, this was progressed into a conditioning program to build strength and endurance within these functional tasks.

3. Functional Integration
The exercise program in stage 2 was functionally integrated, specific to each persons nominated pain provocative activities and directed at their functional goals. The aim was to restore normal functional movement capacity, enhanced body awareness, reduce avoidance, pain behaviours and fear by means of pain control and confrontation in daily life.

Here each participant nominated activities of daily living that provoked their pain and these were rehearsed with the therapist, so that they were confident and mindful of normalising their movement behaviours whilst performing these tasks in activities of daily living. If the participant reported activities they avoided, these were also rehearsed and they were encouraged to confront these activities (without protective behaviours) and include them in their daily activities.

4. Physical activity levels
Patients were asked to carry out some form of physical exercise (such as walking or biking) 3-5 times a week if they weren’t previously doing so (the duration for this was initially based on the patients exercise tolerance and gradually increased to 20-40 minutes duration). They were encouraged to be mindful of how they performed this to facilitate normal movement in a relaxed manner. The type of the physical activity was directed both by the movement classification and the patient’s preference. For example, if the patient had an extension ‘control impairment’ disorder they were first asked to use an exercise bike or bicycle to until they had acquired sufficient control, conditioning and confidence to begin a graduated walking program.

Örebro Musculoskeletal Pain Questionnaire (ÖMPQ)

Linton and Boersma 2003¹

1. Name _____ Phone _____ Date _____

2. Date of Injury _____ Date of birth _____

3. Male Female

4. Were you born in Australia*? Yes No

These questions and statements apply if you have aches or pains, such as back, shoulder or neck pain. Please read and answer questions carefully. Do not take long to answer the questions, however it is important that you answer every question. There is **always** a response for your particular situation.

<p>5. Where do you have pain? Place a tick (✓) for all appropriate sites.</p> <p><input type="checkbox"/> Neck <input type="checkbox"/> Shoulder <input type="checkbox"/> Arm <input type="checkbox"/> Upper Back</p> <p><input type="checkbox"/> Lower Back <input type="checkbox"/> Leg <input type="checkbox"/> Other (state)</p>	2x (max 10)
<p>6. How many days of work have you missed because of pain during the past 18 months? Tick (✓) one.</p> <p><input type="checkbox"/> 0 days (1) <input type="checkbox"/> 1-2 days (2) <input type="checkbox"/> 3-7 days (3) <input type="checkbox"/> 8-14 days (4)</p> <p><input type="checkbox"/> 15-30 days (5) <input type="checkbox"/> 1 month (6) <input type="checkbox"/> 2 months (7) <input type="checkbox"/> 3-6 months (8)</p> <p><input type="checkbox"/> 6-12 months (9) <input type="checkbox"/> over 1 year (10)</p>	
<p>7. How long have you had your current pain problem? Tick (✓) one.</p> <p><input type="checkbox"/> 0-1 week (1) <input type="checkbox"/> 1-2 weeks (2) <input type="checkbox"/> 3-4 weeks (3) <input type="checkbox"/> 4-5 weeks (4)</p> <p><input type="checkbox"/> 6-8 weeks (5) <input type="checkbox"/> 9-11 weeks (6) <input type="checkbox"/> 3-6 months (7) <input type="checkbox"/> 6-9 months (8)</p> <p><input type="checkbox"/> 9-12 months (9) <input type="checkbox"/> over 1 year (10)</p>	
<p>8. Is your work heavy or monotonous? Circle the best alternative.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Not at all Extremely</p>	
<p>9. How would you rate the pain that you have had during the past week? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>No pain Pain as bad as it could be</p>	

* Modified for use by WorkCover NSW (with permission)

¹ Linton SJ, Boersma K. Early identification of patients at risk of developing a persistent back problem: the predictive validity of the Örebro Musculoskeletal Pain Questionnaire. Clin J Pain 2003;19: 80-86.

<p>10. In the past three months, on average, how bad was your pain on a 0-10 scale? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>No pain Pain as bad as it could be</p>	
<p>11. How often would you say that you have experience pain episodes, on average, during the past three months? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Never Always</p>	
<p>12. Based on all things you do to cope, or deal with your pain, on an average day, how much are you able to decrease it? Circle the appropriate number.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Can't decrease it at all Can decrease it completely</p>	10 - x
<p>13. How tense or anxious have you felt in the past week? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Absolutely clam and relaxed As tense and anxious as I've ever felt</p>	
<p>14. How much have you been bothered by feeling depressed in the past week? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Not at all Extremely</p>	
<p>15. In your view, how large is the risk that your current pain may become persistent? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>No risk Very large risk</p>	
<p>16. In your estimation, what are the chances that you will be able to work in six months? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>No chance Very large chance</p>	10 - x
<p>17. If you take into consideration your work routines, management, salary, promotion possibilities and work mates, how satisfied are you with your job? Circle one.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Not satisfied at all Completely satisfied</p>	10 - x

<p>Here are some of the things that other people have told us about their pain. For each statement, circle one number from 0 to 10 to say how much physical activities, such as bending, lifting, walking or driving, would affect your pain.</p>	
<p>18. Physical activity makes my pain worse.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Completely disagree Completely agree</p>	
<p>19. An increase in pain is an indication that I should stop what I'm doing until the pain decreases.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Completely disagree Completely agree</p>	
<p>20. I should not do my normal work with my present pain.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Completely disagree Completely agree</p>	
<p>Here is a list of five activities. Circle the one number that best describes your current ability to participate in each of these activities.</p>	
<p>21. I can do light work for an hour.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Can't do it because of pain problem Can do it without pain being a problem</p>	10 - x
<p>22. I can walk for an hour.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Can't do it because of pain problem Can do it without pain being a problem</p>	10 - x
<p>23. I can do ordinary household chores.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Can't do it because of pain problem Can do it without pain being a problem</p>	10 - x
<p>24. I can do the weekly shopping.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Can't do it because of pain problem Can do it without pain being a problem</p>	10 - x
<p>25. I can sleep at night.</p> <p>0 1 2 3 4 5 6 7 8 9 10</p> <p>Can't do it because of pain problem Can do it without pain being a problem</p>	10 - x

Explanatory Notes

The Örebro Musculoskeletal Pain Questionnaire (ÖMPQ) is a 'yellow flag' screening tool that predicts long-term disability and failure to return to work when completed four to 12 weeks following a soft tissue injury². A cut-off score of 105 has been found to predict those who will recover (with 95 per cent accuracy), those who will have no further sick leave in the next six months (with 81 per cent accuracy), and those who will have long-term sick leave (with 67 per cent accuracy)¹.

The ÖMPQ predicted failure to return to work six months after compensable musculoskeletal injury in a NSW population of workers. The injuries in the study group were mixed, and the ÖMPQ was found to be more specific and sensitive for back injuries. In workers with back injuries screened at four to 12 weeks, a cut-off score of 130 correctly predicted 86 per cent of those who failed to return to work³.

Identification, through the ÖMPQ, of workers at risk of failing to return to work due to personal and environmental factors provides the opportunity for treating practitioners to apply appropriate interventions (including the use of activity programs based on cognitive behavioural strategies) to reduce the risk of long-term disability in injured workers. Evidence indicates that these factors can be changed if they are addressed⁴.

Administering the questionnaire

The ÖMPQ is designed to be a self administered tool completed by the worker in a quiet environment without assistance from any other person. A detailed explanation is provided by the person administering the questionnaire:

"Information from this questionnaire helps us understand your problem better, and it especially helps us evaluate the possible long-term consequences your pain may have. It is important that you read each question carefully and answer it as best you can. There are no right or wrong answers. Please answer every question. If you have difficulty, select the answer that best describes your situation".

Where uncertainty or a request for more information is expressed, encouragement is provided to "answer as best you can". The questionnaire item may be read aloud to assist, however the question should not be rephrased. All questions should be answered, as missing values will reduce validity⁵.

Scoring instructions

- For question 5, count the number of pain sites and multiply by two – this is the score (maximum score allowable is 10).
- For questions 6 and 7 the score is the number bracketed after the ticked box.
- For questions 8, 9, 10, 11, 13, 14, 15, 18, 19 and 20 the score is the number that has been ticked or circled.
- For questions 12, 16, 17, 21, 22, 23, 24 and 25 the score is 10 minus the number that has been circled.
- Write the score in the shaded area beside each item.
- Add up the scores for questions 5 to 25 – this is the total ÖMPQ score.

1 Linton SJ, Boersma K. Early identification of patients at risk of developing a persistent back problem: the predictive validity of the Örebro Musculoskeletal Pain Questionnaire. *Clin J Pain* 2003;19: 80-86.

2 Linton SJ, Hallden K. Can we screen for problematic back pain? A screening questionnaire for predicting outcome in acute and subacute back pain. *Clin J Pain* 1998; 3: 209-215.

3 Dunstan DA, Covic T, Tyson GA, Lennie, IG (2005) Does the ÖMPQ predict outcomes following a work related compensable injury? *International Journal of Rehabilitation Research* 28(4), 369-370.

4 Linton SJ, Ryberg M. A cognitive-behavioral group intervention as prevention for persistent neck and back pain in a non-patient population: a randomized controlled trial. *Pain* 2001; 83-90.

van den Hout JHC, Vlaeyen WS, Heuts PHTG, Zijlema JHL, Wijnen AG. Secondary Prevention of Work-Related Disability in Nonspecific Low Back Pain: Problem-Solving Therapy Help? A Randomized Clinical Trial. *Clinical Journal of Pain* 2003; 19: 87-96.

Marhold C, Linton SJ, Melin L. A cognitive-behavioral return to work program: effects on pain patients with a history of long-term versus short-term sick leave. *Pain* 2001; 91:155-163.

5 Linton SJ. *Understanding pain for better clinical practice – a psychological perspective*. Edinburgh: Elsevier, 2005.

Disclaimer

This publication may contain occupational health and safety and workers compensation information. It may include some of your obligations under the various legislations that WorkCover NSW administers. To ensure you comply with your legal obligations you must refer to the appropriate legislation.

Information on the latest laws can be checked by visiting the NSW legislation website www.legislation.nsw.gov.au.

This publication does not represent a comprehensive statement of the law as it applies to particular problems or to individuals or as a substitute for legal advice. You should seek independent legal advice if you need assistance on the application of the law to your situation.



Catalogue No. **WC05537** WorkCover Publications Hotline **1300 799 003**

WorkCover NSW 92-100 Donnison St Gosford NSW 2250
Locked Bag 2906 Lisarow NSW 2252 WorkCover Assistance Service **13 10 50**
Website **www.workcover.nsw.gov.au**

© Copyright WorkCover NSW 0710